

Table 4B
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
County	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
Adams	799	84.9	0.4	12.3	0.0	0.1	2.4	0.0
Allen	11,288	39.0	25.7	19.9	0.0	6.5	8.9	0.0
Bartholome	1,828	72.6	1.3	21.4	0.0	0.4	4.2	0.0
Benton	235	79.1	0.9	17.0	0.0	0.0	3.0	0.0
Blackford	411	91.2	0.7	3.9	0.2	0.2	3.6	0.0
Boone	736	91.6	1.6	4.6	0.0	0.4	1.8	0.0
Brown	274	92.0	0.7	4.0	0.0	0.0	3.3	0.0
Carroll	348	82.2	0.3	15.2	0.0	0.0	2.3	0.0
Cass	1,557	56.0	0.9	39.2	0.0	1.0	2.8	0.0
Clark	2,905	58.7	10.5	21.8	0.1	0.9	8.1	0.0
Clay	914	91.6	0.3	4.2	0.0	0.5	3.4	0.0
Clinton	1,217	51.6	0.5	46.1	0.1	0.0	1.7	0.0
Crawford	296	96.3	0.0	2.7	0.0	0.0	1.0	0.0
Daviess	816	70.2	0.1	26.2	0.0	0.2	3.2	0.0
Dearborn	1,131	95.0	1.0	1.1	0.1	0.0	2.7	0.0
Decatur	810	96.7	0.1	2.7	0.1	0.0	0.4	0.0
DeKalb	1,106	88.4	0.5	7.2	0.1	0.4	3.3	0.0
Delaware	2,887	76.7	11.2	3.5	0.1	0.6	8.0	0.0
Dubois	1,017	65.0	0.4	30.9	0.1	0.6	3.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4B
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Indiana

Comparison of Racial and Ethnic Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
County	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
Elkhart	7,636	42.6	10.1	41.4	0.0	0.5	5.4	0.0
Fayette	750	95.6	0.8	2.0	0.0	0.3	1.3	0.0
Floyd	1,709	65.5	13.9	9.0	0.1	3.9	7.7	0.0
Fountain	414	94.4	0.7	4.6	0.0	0.2	0.0	0.0
Franklin	367	98.4	0.0	0.5	0.0	0.0	1.1	0.0
Fulton	586	85.7	0.7	11.9	0.0	0.0	1.7	0.0
Gibson	687	84.6	3.5	3.3	0.1	0.0	8.4	0.0
Grant	2,285	66.7	11.1	11.9	0.0	0.2	10.1	0.0
Greene	679	94.6	0.3	3.8	0.0	0.1	1.2	0.0
Hamilton	2,732	73.9	8.2	13.2	0.3	2.0	2.5	0.0
Hancock	1,001	88.5	2.0	4.6	0.2	0.0	4.7	0.0
Harrison	898	91.5	0.1	6.2	0.1	0.7	1.3	0.0
Hendricks	1,447	83.8	6.4	5.1	0.1	2.8	1.7	0.0
Henry	1,286	89.7	1.1	3.8	0.1	0.4	5.0	0.0
Howard	2,898	74.9	10.1	6.7	0.0	0.2	8.1	0.0
Huntington	875	91.4	0.0	3.9	0.0	0.1	4.6	0.0
Jackson	1,427	77.0	1.7	21.0	0.1	0.1	0.1	0.0
Jasper	838	80.7	1.9	16.0	0.0	0.2	1.2	0.0
Jay	588	84.5	0.2	13.1	0.0	0.0	2.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4B
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
County	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
Jefferson	864	82.1	0.8	11.0	0.3	0.6	5.2	0.0
Jennings	1,155	93.5	1.6	4.0	0.1	0.1	0.8	0.0
Johnson	2,558	81.2	1.8	8.1	0.0	4.1	4.8	0.0
Knox	908	88.5	1.9	5.2	0.1	0.0	4.3	0.0
Kosciusko	1,727	66.2	1.0	29.9	0.1	0.5	2.4	0.0
LaGrange	610	77.7	0.7	16.6	0.0	1.1	3.9	0.0
Lake	15,750	24.6	42.2	29.5	0.0	0.3	3.2	0.0
LaPorte	3,248	60.1	20.2	13.9	0.0	0.4	5.3	0.0
Lawrence	1,084	94.7	0.6	2.6	0.0	0.8	1.3	0.0
Madison	3,246	68.1	12.1	12.9	0.1	0.1	6.7	0.0
Marion	31,659	27.1	37.7	26.9	0.1	4.1	4.0	0.0
Marshall	1,329	64.9	0.6	30.6	0.0	0.2	3.8	0.0
Martin	249	94.8	1.2	0.8	0.0	0.4	2.8	0.0
Miami	915	83.1	2.7	5.6	0.4	0.3	7.8	0.1
Monroe	2,512	73.9	5.7	7.1	0.0	4.3	9.0	0.0
Montgomery	1,064	81.1	1.2	16.0	0.0	0.1	1.6	0.0
Morgan	1,450	92.8	0.1	3.4	0.0	0.1	3.5	0.0
Newton	237	86.9	0.4	11.0	0.0	0.4	1.3	0.0
Noble	1,117	67.6	0.3	28.9	0.0	1.0	2.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4B
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
County	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
Ohio	38	*	*	*	*	*	*	*
Orange	669	91.2	2.2	2.8	0.1	0.1	3.4	0.0
Owen	489	93.5	0.2	3.3	0.0	0.4	2.7	0.0
Parke	304	94.7	0.0	4.9	0.0	0.0	0.3	0.0
Perry	434	94.5	0.0	1.4	0.0	0.2	3.9	0.0
Pike	264	95.5	0.0	3.0	0.0	0.0	1.5	0.0
Porter	2,706	70.4	7.6	18.5	0.1	0.6	2.9	0.0
Posey	390	85.9	2.8	3.1	0.0	0.3	7.9	0.0
Pulaski	309	90.9	0.6	5.5	0.0	0.3	2.6	0.0
Putnam	881	93.1	0.7	2.6	0.0	0.2	3.4	0.0
Randolph	698	80.5	0.6	13.8	0.0	0.0	5.2	0.0
Ripley	765	93.2	0.0	4.2	0.0	0.0	2.6	0.0
Rush	333	91.0	0.0	4.2	0.0	0.9	3.9	0.0
Scott	981	93.7	0.4	3.9	0.0	0.1	1.9	0.0
Shelby	1,151	83.1	1.0	14.0	0.0	0.0	1.9	0.0
Spencer	374	83.2	0.5	12.0	0.0	0.0	4.3	0.0
St Joseph	7,045	40.3	27.6	22.7	0.1	1.2	8.2	0.0
Starke	774	91.7	0.1	6.5	0.0	0.0	1.7	0.0
Steuben	1,246	87.6	0.6	8.6	0.0	1.0	2.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4B
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
County	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
Sullivan	507	94.3	0.8	3.2	0.0	0.0	1.8	0.0
Switzerland	179	96.6	0.6	1.7	0.0	0.0	1.1	0.0
Tippecanoe	4,738	53.3	10.4	25.8	0.2	5.0	5.3	0.0
Tipton	260	89.2	0.4	5.8	0.4	0.0	4.2	0.0
Union	190	94.2	0.0	1.6	0.0	0.0	4.2	0.0
Vanderburg	3,836	66.2	19.7	6.6	0.1	0.8	6.6	0.0
Vermillion	368	96.7	0.0	1.9	0.0	0.3	1.1	0.0
Vigo	2,661	75.9	8.5	4.9	0.0	0.8	9.8	0.0
Wabash	922	84.3	0.9	7.6	0.0	0.7	6.6	0.0
Warren	278	98.6	0.0	1.4	0.0	0.0	0.0	0.0
Warrick	956	88.7	1.8	4.3	0.0	0.2	5.0	0.0
Washington	702	93.7	0.0	3.4	0.0	0.1	2.7	0.0
Wayne	2,232	71.6	6.0	9.8	0.1	0.3	12.2	0.0
Wells	567	86.1	0.7	7.6	0.0	0.0	5.6	0.0
White	831	74.5	0.1	23.5	0.0	0.2	1.7	0.0
Whitley	746	93.8	0.4	3.9	0.0	0.1	1.7	0.0
Unknown Counties	5	*	*	*	*	*	*	*
Indiana	171,189	57.4	17.2	18.6	0.1	1.8	4.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4B
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Racial and Ethnic Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
County	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
Nation (Prior Year)	8,672,965	32.5	19.1	39.7	0.9	2.5	2.7	2.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5B
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution				
County	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Adams	799	37.4	18.1	44.4
Allen	11,288	32.7	20.3	47.0
Bartholome	1,828	41.6	17.8	40.6
Benton	235	27.7	17.4	54.9
Blackford	411	34.5	21.4	44.0
Boone	736	36.0	19.4	44.6
Brown	274	36.1	15.7	48.2
Carroll	348	34.5	16.1	49.4
Cass	1,557	35.6	17.0	47.4
Clark	2,905	34.4	21.1	44.5
Clay	914	36.7	17.6	45.7
Clinton	1,217	34.4	17.9	47.7
Crawford	296	36.5	16.2	47.3
Daviess	816	32.6	19.0	48.4
Dearborn	1,131	41.2	16.9	41.9
Decatur	810	36.3	17.0	46.7
DeKalb	1,106	34.1	24.0	42.0
Delaware	2,887	41.0	18.6	40.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5B
2011 Pediatric Nutrition Surveillance ⁽¹⁾
Indiana
Comparison of Age Distribution by County
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution				
County	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Dubois	1,017	38.0	15.9	46.1
Elkhart	7,636	33.7	20.0	46.3
Fayette	750	34.8	22.0	43.2
Floyd	1,709	37.4	21.3	41.3
Fountain	414	36.5	18.4	45.2
Franklin	367	35.1	23.7	41.1
Fulton	586	32.3	21.5	46.2
Gibson	687	44.1	19.8	36.1
Grant	2,285	36.1	17.8	46.1
Greene	679	34.3	19.7	45.9
Hamilton	2,732	38.9	18.4	42.7
Hancock	1,001	42.0	20.3	37.8
Harrison	898	38.2	17.3	44.5
Hendricks	1,447	38.5	21.1	40.4
Henry	1,286	34.8	19.8	45.4
Howard	2,898	36.7	18.5	44.8
Huntington	875	37.8	20.0	42.2
Jackson	1,427	36.0	17.3	46.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5B
2011 Pediatric Nutrition Surveillance (1)
Indiana
Comparison of Age Distribution by County
Children Aged < 5 Years (2)

Comparison of Age Distribution				
County	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Jasper	838	38.2	20.5	41.3
Jay	588	37.4	16.7	45.9
Jefferson	864	38.9	18.1	43.1
Jennings	1,155	33.0	18.6	48.4
Johnson	2,558	42.8	17.1	40.0
Knox	908	31.7	23.2	45.0
Kosciusko	1,727	35.5	19.9	44.6
LaGrange	610	33.9	17.0	49.0
Lake	15,750	37.7	18.3	44.0
LaPorte	3,248	38.9	17.5	43.6
Lawrence	1,084	39.6	19.4	41.1
Madison	3,246	42.9	17.9	39.2
Marion	31,659	34.3	23.7	42.0
Marshall	1,329	38.8	17.0	44.2
Martin	249	38.2	16.5	45.4
Miami	915	35.8	16.8	47.3
Monroe	2,512	39.0	18.4	42.6
Montgomery	1,064	37.0	20.0	43.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5B
2011 Pediatric Nutrition Surveillance (1)
Indiana
Comparison of Age Distribution by County
Children Aged < 5 Years (2)

Comparison of Age Distribution				
County	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Morgan	1,450	44.7	14.8	40.5
Newton	237	38.4	14.3	47.3
Noble	1,117	30.9	23.5	45.6
Ohio	38	*	*	*
Orange	669	36.9	15.8	47.2
Owen	489	33.1	21.3	45.6
Parke	304	36.8	18.8	44.4
Perry	434	47.7	15.2	37.1
Pike	264	45.1	15.5	39.4
Porter	2,706	41.4	18.0	40.6
Posey	390	44.6	19.0	36.4
Pulaski	309	39.2	17.8	43.0
Putnam	881	34.6	20.5	44.8
Randolph	698	35.8	20.8	43.4
Ripley	765	35.0	20.7	44.3
Rush	333	36.9	19.5	43.5
Scott	981	33.7	19.2	47.1
Shelby	1,151	41.4	18.9	39.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5B
2011 Pediatric Nutrition Surveillance (1)
Indiana
Comparison of Age Distribution by County
Children Aged < 5 Years (2)

Comparison of Age Distribution				
County	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Spencer	374	38.5	17.9	43.6
St Joseph	7,045	35.6	21.3	43.1
Starke	774	34.1	18.1	47.8
Steuben	1,246	31.0	18.5	50.5
Sullivan	507	46.2	18.5	35.3
Switzerlan	179	33.0	18.4	48.6
Tippecanoe	4,738	35.7	20.2	44.2
Tipton	260	35.4	16.9	47.7
Union	190	34.2	21.6	44.2
Vanderburg	3,836	46.0	17.3	36.7
Vermillion	368	36.4	20.7	42.9
Vigo	2,661	40.5	20.4	39.0
Wabash	922	34.6	16.5	48.9
Warren	278	34.5	18.7	46.8
Warrick	956	41.8	18.6	39.5
Washington	702	34.3	18.8	46.9
Wayne	2,232	35.1	21.1	43.9
Wells	567	37.4	18.5	44.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5B
2011 Pediatric Nutrition Surveillance (1)
Indiana
Comparison of Age Distribution by County
Children Aged < 5 Years (2)

Comparison of Age Distribution				
County	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
White	831	35.9	18.2	46.0
Whitley	746	33.6	18.0	48.4
Unknown Counties	5	*	*	*
Indiana	171,189	36.6	19.9	43.5
Nation (Prior Year)	8,672,965	33.7	22.1	44.2

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under-weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Adams	186	8.6 (39)	6.5 (39)	789	8.6 (85)	2.4 (55)	10.8 (39)	355	20.0 (78)	10.7 (9)	536	22.0 (87)
Allen	2,586	10.0 (55)	4.6 (7)	11,131	6.4 (42)	2.4 (55)	11.7 (58)	5,251	16.4 (38)	13.8 (27)	7,956	22.4 (88)
Bartholome	502	8.8 (42)	5.6 (19)	1,733	6.8 (54)	1.6 (26)	10.0 (20)	733	16.2 (35)	14.2 (33)	1,132	9.4 (29)
Benton	45	* *	* *	233	4.7 (8)	0.0 (1)	12.4 (71)	125	20.0 (78)	20.0 (86)	169	8.3 (20)
Blackford	96	* *	* *	410	6.3 (37)	3.4 (80)	11.2 (49)	179	10.6 (2)	15.6 (48)	267	11.2 (41)
Boone	180	6.1 (8)	7.2 (48)	718	7.0 (65)	5.2 (90)	13.6 (86)	320	19.1 (75)	18.8 (78)	478	5.0 (4)
Brown	60	* *	* *	268	5.2 (13)	0.4 (4)	9.3 (14)	128	14.1 (9)	14.1 (32)	181	2.8 (1)
Carroll	80	* *	* *	339	6.5 (43)	1.5 (21)	11.2 (49)	173	17.3 (57)	14.5 (35)	240	18.3 (76)
Cass	349	6.3 (10)	6.6 (41)	1,520	4.7 (8)	3.3 (76)	12.2 (67)	720	15.4 (26)	16.7 (63)	1,072	10.2 (36)
Clark	682	7.9 (26)	6.0 (30)	2,867	6.2 (34)	1.5 (21)	12.3 (68)	1,271	17.9 (62)	16.5 (61)	1,998	11.9 (44)
Clay	213	6.1 (8)	4.7 (8)	902	7.6 (76)	1.3 (16)	12.0 (64)	407	16.7 (42)	16.7 (63)	623	13.8 (56)

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this county's rate to other counties. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
 - (7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under-weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Clinton	259	6.9 (11)	8.1 (61)	1,188	7.2 (69)	1.6 (26)	13.0 (77)	569	19.0 (74)	16.5 (61)	847	12.5 (46)
Crawford	74	* *	* *	287	6.6 (49)	0.3 (3)	9.8 (19)	136	18.4 (67)	14.7 (38)	195	8.7 (24)
Daviess	206	7.8 (24)	2.9 (3)	802	6.7 (50)	3.9 (84)	12.1 (66)	383	15.7 (29)	19.8 (85)	536	17.2 (72)
Dearborn	308	10.4 (59)	8.4 (66)	1,117	6.5 (43)	1.3 (16)	10.0 (20)	463	16.8 (46)	13.6 (25)	764	11.6 (43)
Decatur	193	12.4 (72)	6.2 (35)	795	6.8 (54)	1.6 (26)	12.7 (73)	374	16.3 (37)	17.9 (71)	566	8.5 (23)
DeKalb	272	8.5 (37)	6.6 (41)	1,086	5.5 (16)	2.4 (55)	11.7 (58)	460	17.0 (50)	17.2 (66)	757	18.1 (75)
Delaware	816	10.7 (61)	4.3 (6)	2,828	7.1 (66)	4.9 (89)	10.6 (35)	1,140	16.7 (42)	15.4 (46)	1,850	22.7 (89)
Dubois	242	4.5 (3)	5.8 (25)	992	5.4 (15)	2.3 (51)	10.3 (31)	455	15.8 (31)	14.5 (35)	677	8.7 (24)
Elkhart	1,769	8.1 (32)	8.2 (62)	7,467	5.5 (16)	1.5 (21)	11.0 (45)	3,487	16.9 (49)	15.9 (52)	5,244	6.5 (9)
Fayette	181	7.7 (20)	8.3 (65)	740	7.7 (77)	2.7 (68)	11.1 (47)	319	13.5 (6)	14.7 (38)	496	18.3 (76)
Floyd	431	9.5 (52)	6.3 (37)	1,644	7.4 (72)	2.1 (43)	12.7 (73)	696	20.8 (83)	17.2 (66)	1,133	13.1 (50)

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this county's rate to other counties. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
 - (7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			(8)	
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Fountain	100	12.0 (69)	6.0 (30)	406	6.9 (60)	1.2 (12)	7.6 (3)	181	18.8 (71)	12.7 (19)	283	13.1 (50)
Franklin	89	* *	* *	362	6.9 (60)	1.1 (8)	11.9 (63)	149	22.1 (87)	19.5 (84)	240	21.7 (85)
Fulton	115	4.3 (2)	6.1 (32)	575	3.3 (1)	3.5 (82)	12.3 (68)	266	14.7 (18)	16.2 (55)	429	14.2 (59)
Gibson	187	7.0 (13)	5.3 (12)	670	6.7 (50)	2.5 (64)	12.8 (76)	251	17.1 (53)	20.3 (87)	437	14.6 (63)
Grant	528	10.2 (56)	5.5 (15)	2,225	6.7 (50)	2.4 (55)	11.4 (52)	1,038	16.0 (33)	14.0 (30)	1,539	19.4 (81)
Greene	165	9.1 (48)	5.5 (15)	672	6.1 (29)	1.6 (26)	13.1 (80)	312	22.8 (89)	18.9 (80)	448	8.9 (27)
Hamilton	645	9.1 (48)	6.5 (39)	2,683	6.2 (34)	2.4 (55)	7.7 (5)	1,154	14.6 (13)	9.4 (5)	1,822	15.7 (67)
Hancock	275	6.9 (11)	8.7 (69)	982	9.2 (91)	1.7 (31)	13.0 (77)	373	15.3 (23)	15.8 (51)	641	12.9 (49)
Harrison	235	7.2 (14)	6.8 (44)	868	6.3 (37)	2.4 (55)	11.4 (52)	388	16.5 (39)	17.5 (69)	591	3.9 (2)
Hendricks	434	9.0 (46)	5.8 (25)	1,409	7.4 (72)	4.0 (85)	9.3 (14)	578	16.8 (46)	12.3 (15)	844	21.0 (84)
Henry	289	7.6 (19)	7.3 (49)	1,269	6.0 (27)	3.5 (82)	10.2 (25)	579	20.2 (80)	15.2 (45)	912	11.0 (40)

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this county's rate to other counties. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
 - (7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under-weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Howard	651	8.0 (29)	4.9 (9)	2,810	6.5 (43)	2.3 (51)	11.7 (58)	1,267	16.7 (42)	15.7 (50)	2,041	21.8 (86)
Huntington	215	7.9 (26)	5.1 (10)	870	7.1 (66)	2.4 (55)	8.6 (9)	363	15.4 (26)	10.2 (7)	585	8.7 (24)
Jackson	328	8.8 (42)	8.2 (62)	1,409	6.5 (43)	0.9 (7)	11.6 (57)	652	19.3 (76)	17.0 (65)	982	5.3 (6)
Jasper	203	5.9 (7)	9.9 (70)	810	6.3 (37)	2.3 (51)	13.0 (77)	345	18.0 (64)	15.9 (52)	555	18.7 (79)
Jay	146	10.3 (58)	2.7 (2)	581	6.9 (60)	1.2 (12)	11.4 (52)	271	16.2 (35)	15.5 (47)	391	11.5 (42)
Jefferson	204	12.3 (71)	7.4 (52)	849	7.4 (72)	1.2 (12)	13.4 (84)	353	21.0 (84)	21.2 (89)	596	8.2 (19)
Jennings	262	9.5 (52)	6.9 (45)	1,137	7.7 (77)	3.3 (76)	13.6 (86)	545	17.6 (59)	18.0 (72)	802	16.6 (71)
Johnson	708	8.3 (36)	7.9 (59)	2,484	5.6 (20)	2.7 (68)	6.8 (2)	1,016	14.6 (13)	10.3 (8)	1,633	12.6 (48)
Knox	240	10.8 (63)	3.8 (4)	890	7.9 (80)	3.4 (80)	10.8 (39)	404	15.3 (23)	15.6 (48)	544	6.6 (10)
Kosciusko	415	7.7 (20)	6.3 (37)	1,698	5.8 (24)	1.2 (12)	12.6 (72)	758	18.3 (65)	18.2 (75)	1,189	7.3 (11)
LaGrange	130	11.5 (67)	6.9 (45)	596	5.2 (13)	2.2 (44)	11.4 (52)	293	17.4 (58)	16.0 (54)	401	6.0 (8)

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this county's rate to other counties. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
 - (7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under-weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Lake	3,739	9.6 (54)	5.9 (28)	15,387	6.3 (37)	2.2 (44)	10.2 (25)	6,834	14.8 (19)	12.2 (14)	10,556	15.3 (65)
LaPorte	776	9.3 (50)	5.5 (15)	3,160	6.1 (29)	3.1 (74)	11.2 (49)	1,400	16.8 (46)	13.6 (25)	2,139	14.7 (64)
Lawrence	263	10.6 (60)	5.7 (23)	1,067	6.8 (54)	1.3 (16)	11.1 (47)	446	16.1 (34)	12.6 (17)	712	9.7 (33)
Madison	897	8.6 (39)	5.7 (23)	3,171	8.9 (87)	2.4 (55)	10.2 (25)	1,253	16.5 (39)	14.2 (33)	2,144	28.5 (90)
Marion	8,803	9.3 (50)	6.1 (32)	30,955	6.2 (34)	4.2 (86)	10.7 (37)	13,110	15.3 (23)	14.0 (30)	19,568	17.4 (73)
Marshall	354	4.2 (1)	7.6 (55)	1,300	7.2 (69)	1.1 (8)	13.1 (80)	581	19.4 (77)	16.2 (55)	874	7.7 (14)
Martin	61	* *	* *	245	5.7 (22)	0.4 (4)	12.7 (73)	110	12.7 (4)	16.4 (59)	152	5.9 (7)
Miami	199	9.0 (46)	5.5 (15)	897	7.1 (66)	1.6 (26)	9.0 (12)	432	14.6 (13)	11.8 (12)	620	11.9 (44)
Monroe	588	8.8 (42)	10.0 (71)	2,462	7.3 (71)	1.7 (31)	10.4 (33)	1,060	18.6 (68)	14.6 (37)	1,691	13.7 (55)
Montgomery	271	7.7 (20)	11.1 (72)	1,049	6.7 (50)	3.1 (74)	9.7 (18)	452	17.9 (62)	14.8 (40)	723	13.6 (54)
Morgan	403	8.7 (41)	8.4 (66)	1,402	6.8 (54)	1.9 (37)	10.7 (37)	577	18.7 (70)	18.0 (72)	929	13.3 (53)

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this county's rate to other counties. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
 - (7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under-weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Newton	57	* *	* *	231	4.3 (3)	7.4 (91)	13.9 (88)	108	15.7 (29)	12.0 (13)	146	4.1 (3)
Noble	285	8.1 (32)	5.6 (19)	1,082	8.0 (81)	2.6 (67)	11.8 (62)	506	14.6 (13)	13.8 (27)	727	19.3 (80)
Ohio	7	* *	* *	37	* *	* *	* *	21	* *	* *	27	* *
Orange	149	8.1 (32)	1.3 (1)	653	5.8 (24)	2.5 (64)	9.0 (12)	310	14.8 (19)	14.8 (40)	454	10.1 (35)
Owen	107	5.6 (4)	8.4 (66)	478	6.5 (43)	1.7 (31)	13.2 (82)	222	13.5 (6)	18.5 (77)	340	7.4 (12)
Parke	74	* *	* *	301	6.0 (27)	3.3 (76)	10.0 (20)	134	13.4 (5)	13.4 (24)	216	7.4 (12)
Perry	124	14.5 (73)	5.6 (19)	422	9.0 (88)	4.3 (88)	10.2 (25)	162	11.1 (3)	9.3 (4)	293	34.1 (91)
Pike	66	* *	* *	256	7.8 (79)	2.0 (39)	10.9 (44)	100	17.0 (50)	19.0 (83)	166	7.8 (15)
Porter	728	7.8 (24)	7.7 (57)	2,641	7.4 (72)	2.2 (44)	10.5 (34)	1,083	14.3 (11)	12.3 (15)	1,779	13.8 (56)
Posey	104	7.7 (20)	7.7 (57)	382	4.7 (8)	1.8 (36)	8.4 (8)	140	14.3 (11)	7.9 (1)	243	14.4 (62)
Pulaski	73	* *	* *	299	4.3 (3)	2.0 (39)	14.0 (89)	135	20.7 (82)	12.6 (17)	208	12.5 (46)

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this county's rate to other counties. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
 - (7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under-weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Putnam	213	8.9 (45)	6.1 (32)	869	6.1 (29)	2.8 (70)	12.3 (68)	388	18.6 (68)	16.2 (55)	592	8.3 (20)
Randolph	161	5.6 (4)	6.2 (35)	695	6.8 (54)	1.7 (31)	9.6 (17)	306	13.7 (8)	13.1 (22)	475	9.5 (30)
Ripley	169	10.7 (61)	5.9 (28)	755	4.4 (5)	1.7 (31)	10.1 (23)	344	16.6 (41)	14.8 (40)	548	8.0 (18)
Rush	86	* *	* *	331	6.9 (60)	4.2 (86)	5.4 (1)	144	14.6 (13)	8.3 (3)	221	13.1 (50)
Scott	223	10.8 (63)	5.8 (25)	963	6.1 (29)	3.0 (73)	11.7 (58)	445	15.1 (21)	18.0 (72)	664	10.8 (39)
Shelby	307	7.2 (14)	4.2 (5)	1,120	8.7 (86)	0.8 (6)	13.5 (85)	450	17.6 (59)	17.6 (70)	755	14.2 (59)
Spencer	93	* *	* *	361	9.1 (90)	1.4 (19)	16.1 (91)	162	22.2 (88)	21.0 (88)	253	9.5 (30)
St Joseph	1,844	10.2 (56)	7.6 (55)	6,898	5.5 (16)	2.9 (72)	10.2 (25)	3,010	15.6 (28)	13.0 (21)	4,369	14.1 (58)
Starke	163	8.0 (29)	7.4 (52)	748	5.6 (20)	1.1 (8)	14.4 (90)	361	15.8 (31)	18.8 (78)	570	15.3 (65)
Steuben	250	8.0 (29)	11.2 (73)	1,220	4.6 (7)	2.2 (44)	9.3 (14)	626	17.1 (53)	12.8 (20)	900	10.6 (37)
Sullivan	152	7.2 (14)	5.3 (12)	502	9.0 (88)	1.4 (19)	13.3 (83)	180	20.6 (81)	18.9 (80)	330	10.6 (37)

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this county's rate to other counties. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
 - (7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2011 Pediatric Nutrition Surveillance (1)
Indiana
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			Number	%
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Switzerland	32	* *	* *	176	4.5 (6)	2.3 (51)	10.2 (25)	85	* *	* *	133	15.8 (68)
Tippecanoe	1,142	7.4 (17)	7.3 (49)	4,664	4.8 (11)	2.4 (55)	10.6 (35)	2,076	17.2 (55)	13.2 (23)	3,193	14.2 (59)
Tipton	56	* *	* *	253	5.5 (16)	2.0 (39)	10.3 (31)	122	8.2 (1)	14.8 (40)	179	7.8 (15)
Union	44	* *	* *	187	3.7 (2)	1.1 (8)	8.6 (9)	86	* *	* *	123	20.3 (83)
Vanderburg	1,076	10.8 (63)	5.2 (11)	3,727	8.0 (81)	2.2 (44)	10.1 (23)	1,386	18.8 (71)	14.8 (40)	2,428	7.8 (15)
Vermillion	84	* *	* *	366	5.7 (22)	2.2 (44)	12.0 (64)	159	17.0 (50)	18.9 (80)	262	17.9 (74)
Vigo	753	8.5 (37)	6.6 (41)	2,620	6.9 (60)	1.5 (21)	10.8 (39)	1,029	18.3 (65)	16.4 (59)	1,724	9.6 (32)
Wabash	198	5.6 (4)	7.1 (47)	898	5.8 (24)	1.9 (37)	8.0 (6)	443	16.7 (42)	9.5 (6)	617	8.4 (22)
Warren	64	* *	* *	277	5.1 (12)	2.2 (44)	10.8 (39)	129	17.8 (61)	16.3 (58)	186	9.7 (33)
Warrick	233	11.6 (68)	7.3 (49)	931	8.5 (84)	2.8 (70)	8.9 (11)	370	14.1 (9)	11.1 (11)	623	15.9 (69)
Washington	146	8.2 (35)	8.2 (62)	679	6.5 (43)	1.5 (21)	10.8 (39)	320	21.9 (86)	17.2 (66)	480	5.0 (4)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under-weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Wayne	545	7.5 (18)	7.9 (59)	2,201	6.3 (37)	2.0 (39)	11.0 (45)	976	18.9 (73)	13.9 (29)	1,458	18.4 (78)
Wells	143	11.2 (66)	5.6 (19)	558	6.8 (54)	0.2 (2)	8.2 (7)	243	21.0 (84)	10.7 (9)	385	9.1 (28)
White	189	7.9 (26)	7.4 (52)	819	6.1 (29)	3.3 (76)	7.6 (3)	377	15.1 (21)	8.2 (2)	567	20.1 (82)
Whitley	167	12.0 (69)	5.4 (14)	730	8.1 (83)	2.5 (64)	11.5 (56)	349	17.2 (55)	18.3 (76)	498	16.5 (70)
Unknown Counties	2	* *	* *	5	* *	* *	* *	2	* *	* *	3	* *
Indiana	42,957	8.9	6.4	167,541	6.4	2.6	10.8	73,424	16.4	14.3	112,865	14.8
Nation (Prior Year)	1,851,546	9.0	6.3	8,057,124	6.3	3.6	12.4	3,572,939	16.1	14.4	5,391,027	14.6

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Rank compares this county's rate to other counties. Rank 1 = best rate.
- (5) Infants born during the reporting period included in the analysis.
- (6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
- (7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
- (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7B
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
County	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
Adams	186	81.7 (3)	133	23.3 (11)	142	7.0 (43)	109	22.9 (4)	133	9.0 (30)	364	57.7 (84)	768	39.3 (36)
Allen	2,592	64.0 (47)	1,686	18.8 (22)	2,000	8.7 (31)	1,228	14.4 (28)	1,686	8.5 (32)	5,189	70.8 (67)	10,949	32.4 (25)
Bartholome	502	68.5 (36)	319	17.9 (24)	276	10.1 (23)	273	12.8 (32)	319	8.2 (36)	763	86.8 (9)	1,732	48.3 (59)
Benton	45	* *	30	* *	31	* *	22	* *	30	* *	132	69.7 (70)	230	45.2 (49)
Blackford	96	* *	61	* *	76	* *	58	* *	61	* *	184	75.5 (43)	391	52.7 (72)
Boone	180	55.6 (67)	102	16.7 (28)	113	6.2 (54)	68	* *	102	13.7 (6)	335	74.9 (46)	717	48.7 (61)
Brown	60	* *	54	* *	46	* *	33	* *	54	* *	134	83.6 (12)	262	45.8 (50)
Carroll	80	* *	63	* *	52	* *	53	* *	63	* *	178	88.8 (6)	330	19.1 (13)
Cass	356	63.2 (49)	309	20.1 (19)	247	6.9 (45)	229	11.4 (43)	309	5.5 (55)	764	60.9 (82)	1,505	21.9 (16)
Clark	685	60.9 (57)	456	20.8 (18)	471	10.8 (15)	322	18.6 (11)	456	10.3 (18)	1,349	77.7 (29)	2,851	38.9 (35)
Clay	213	64.8 (44)	161	12.4 (55)	152	5.3 (60)	150	10.7 (47)	161	5.6 (54)	433	80.1 (23)	882	54.4 (77)
Clinton	259	71.4 (26)	259	11.2 (59)	196	10.2 (20)	208	9.1 (51)	259	3.5 (66)	595	64.9 (74)	1,169	8.2 (2)
Crawford	74	* *	61	* *	45	* *	52	* *	61	* *	145	82.8 (15)	286	51.4 (68)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7B
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
County	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
Daviess	206	58.7 (62)	97	* *	124	7.3 (41)	71	* *	97	* *	409	68.7 (71)	803	40.1 (37)
Dearborn	311	63.7 (48)	199	13.6 (47)	179	7.3 (41)	163	14.1 (30)	199	8.5 (32)	485	77.7 (29)	1,077	57.6 (88)
Decatur	193	70.5 (27)	163	14.1 (43)	131	6.9 (45)	138	12.3 (36)	163	11.0 (14)	386	76.7 (37)	779	55.1 (84)
DeKalb	273	86.4 (1)	178	19.7 (20)	225	8.4 (34)	129	20.9 (7)	178	11.8 (10)	483	74.9 (46)	1,096	49.2 (64)
Delaware	817	59.0 (61)	452	15.3 (38)	426	8.9 (28)	373	12.3 (36)	452	4.4 (57)	1,204	66.9 (72)	2,753	16.3 (8)
Dubois	242	77.3 (6)	226	23.5 (10)	173	13.9 (5)	177	15.8 (23)	226	9.7 (23)	472	75.0 (45)	967	37.3 (30)
Elkhart	1,780	75.0 (14)	1,235	25.3 (6)	1,259	12.4 (9)	909	15.4 (24)	1,235	9.7 (23)	3,669	75.1 (44)	7,337	7.4 (1)
Fayette	181	57.5 (65)	120	14.2 (42)	146	6.2 (54)	76	* *	120	4.2 (59)	333	77.5 (33)	725	53.7 (75)
Floyd	434	64.1 (45)	228	18.9 (21)	292	10.3 (18)	188	12.2 (38)	228	7.5 (41)	739	76.3 (38)	1,651	44.8 (48)
Fountain	100	69.0 (34)	70	* *	65	* *	68	* *	70	* *	195	72.8 (61)	397	54.9 (82)
Franklin	89	* *	61	* *	61	* *	43	* *	61	* *	159	86.2 (10)	351	54.7 (81)
Fulton	115	73.9 (16)	141	14.9 (39)	116	7.8 (38)	89	* *	141	11.3 (13)	276	79.7 (25)	570	41.9 (40)
Gibson	190	51.6 (70)	167	10.8 (60)	103	6.8 (48)	134	9.7 (50)	167	4.2 (59)	259	73.0 (58)	643	50.7 (66)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7B
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
County	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
Grant	530	48.7 (72)	434	11.3 (58)	403	6.5 (50)	325	5.2 (56)	434	4.1 (62)	1,079	75.9 (42)	2,199	16.2 (7)
Greene	165	63.0 (51)	145	7.6 (65)	112	3.6 (64)	118	8.5 (53)	145	4.1 (62)	322	92.5 (2)	655	40.5 (38)
Hamilton	647	77.1 (7)	553	27.7 (2)	451	15.7 (2)	442	21.3 (6)	553	14.8 (5)	1,199	87.1 (8)	2,579	16.8 (9)
Hancock	276	76.4 (10)	172	15.7 (34)	184	10.3 (18)	135	25.9 (1)	172	11.6 (12)	394	73.1 (57)	951	11.3 (5)
Harrison	235	73.2 (18)	178	11.8 (57)	138	8.7 (31)	144	16.7 (17)	178	8.4 (34)	415	89.6 (4)	856	36.2 (27)
Hendricks	436	72.0 (23)	169	21.9 (15)	235	10.6 (16)	158	22.8 (5)	169	11.8 (10)	596	77.7 (29)	1,414	36.4 (28)
Henry	290	69.0 (34)	220	12.7 (54)	232	6.5 (50)	161	14.3 (29)	220	9.1 (29)	606	60.2 (83)	1,232	52.9 (73)
Howard	654	50.6 (71)	601	14.0 (44)	554	6.3 (52)	444	11.5 (42)	601	7.8 (37)	1,332	74.7 (50)	2,780	36.8 (29)
Huntington	216	68.1 (37)	139	15.8 (33)	157	7.6 (40)	108	17.6 (15)	139	10.1 (20)	385	74.8 (48)	843	55.3 (85)
Jackson	331	70.1 (29)	234	16.7 (28)	221	6.3 (52)	179	16.2 (22)	234	9.4 (27)	689	76.3 (38)	1,405	42.7 (43)
Jasper	203	60.6 (58)	174	15.5 (36)	139	12.9 (8)	135	18.5 (12)	174	12.1 (9)	364	47.5 (88)	807	42.8 (44)
Jay	146	63.0 (51)	89	* *	84	* *	84	* *	88	* *	280	53.2 (86)	561	46.3 (52)
Jefferson	208	69.7 (31)	145	13.1 (49)	142	10.6 (16)	122	14.8 (27)	145	7.6 (40)	382	74.6 (51)	825	37.8 (32)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7B
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
County	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
Monroe	589	80.6 (4)	511	30.1 (1)	431	18.1 (1)	390	25.1 (2)	511	20.0 (1)	1,103	88.9 (5)	2,418	18.4 (12)
Montgomery	273	70.0 (30)	182	15.4 (37)	191	5.8 (57)	139	15.1 (26)	182	7.1 (42)	481	64.0 (75)	1,033	43.7 (45)
Morgan	405	72.6 (20)	332	16.3 (32)	203	5.9 (56)	266	17.3 (16)	332	9.6 (25)	607	62.9 (80)	1,340	50.8 (67)
Newton	57	* *	48	* *	44	* *	38	* *	48	* *	113	11.5 (89)	230	48.7 (61)
Noble	286	72.0 (23)	129	24.8 (7)	184	13.0 (7)	95	* *	129	19.4 (2)	528	63.1 (79)	1,110	32.3 (24)
Ohio	7	* *	3	* *	4	* *	3	* *	3	* *	22	* *	36	* *
Orange	149	75.2 (13)	149	18.8 (22)	108	10.2 (20)	111	20.7 (8)	149	6.0 (51)	327	71.6 (65)	634	23.7 (19)
Owen	107	75.7 (11)	79	* *	92	* *	51	* *	79	* *	230	77.0 (34)	473	57.1 (87)
Parke	74	* *	46	* *	51	* *	36	* *	46	* *	138	81.2 (20)	297	51.9 (70)
Perry	124	52.4 (69)	115	5.2 (67)	71	* *	117	7.7 (54)	115	3.5 (66)	166	65.7 (73)	399	59.1 (91)
Pike	68	* *	60	* *	41	* *	58	* *	60	* *	103	74.8 (48)	241	58.5 (90)
Porter	731	72.1 (21)	619	15.7 (34)	440	11.8 (10)	463	12.7 (33)	619	8.4 (34)	1,135	77.9 (28)	2,570	21.6 (15)
Posey	105	53.3 (68)	112	7.1 (66)	58	* *	83	* *	112	6.3 (49)	144	72.9 (60)	364	46.4 (53)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7B
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
County	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)		Smoking in Household					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% (Rank)	Number	% (Rank)				
Pulaski	74	* *	67	* *	45	* *	49	* *	67	* *	137	73.0 (58)	296	48.0 (56)
Putnam	214	65.0 (43)	137	13.1 (49)	151	6.6 (49)	94	* *	137	3.6 (64)	406	73.2 (56)	863	56.3 (86)
Randolph	162	61.7 (55)	124	24.2 (8)	125	8.8 (29)	80	* *	124	12.9 (7)	311	50.2 (87)	666	42.6 (42)
Ripley	169	58.6 (63)	119	5.0 (68)	132	5.3 (60)	85	* *	119	2.5 (68)	348	77.0 (34)	739	54.4 (77)
Rush	88	* *	52	* *	63	* *	38	* *	52	* *	148	77.0 (34)	321	48.0 (56)
Scott	225	47.1 (73)	129	13.2 (48)	140	5.7 (58)	104	16.3 (21)	129	10.9 (16)	477	76.3 (38)	951	54.2 (76)
Shelby	309	67.6 (39)	258	14.0 (44)	189	6.9 (45)	200	13.5 (31)	258	7.8 (37)	475	72.8 (61)	1,093	49.1 (63)
Spencer	93	* *	74	* *	65	* *	75	* *	74	* *	170	91.2 (3)	357	54.9 (82)
St Joseph	1,850	77.0 (8)	910	26.9 (4)	1,195	13.4 (6)	769	15.3 (25)	910	9.3 (28)	3,128	77.7 (29)	6,860	16.8 (9)
Starke	163	69.3 (32)	141	12.8 (53)	127	8.7 (31)	107	11.2 (44)	141	6.4 (47)	385	80.8 (22)	739	51.8 (69)
Steuben	253	82.6 (2)	181	22.1 (13)	203	11.8 (10)	132	19.7 (9)	181	11.0 (14)	655	78.5 (26)	1,213	48.4 (60)
Sullivan	152	71.7 (25)	100	14.0 (44)	79	* *	76	* *	100	9.0 (30)	185	74.1 (54)	493	54.6 (79)
Switzerlan	34	* *	29	* *	32	* *	25	* *	29	* *	85	* *	175	58.3 (89)

(1) Reporting period is January 1 through December 31.
 (2) TV Viewing and Smoking in Household data are not reported by all contributors.
 (3) Analyses based on one record per child.
 (4) Excludes records with unknown data or errors.
 (5) Rank compares this county's rate to other counties. Rank 1 = best rate.
 (6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 (7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
 (8) Children aged 2 years and older included in the analyses.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7B
2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
County	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
Whitley	167	79.6 (5)	111	27.0 (3)	127	15.0 (4)	82	* *	111	18.9 (3)	373	82.8 (15)	724	41.7 (39)
Unknown Counties	2	* *	0	* *	1	* *	1	* *	0	* *	2	* *	5	* *
Indiana	43,169	66.4	28,849	18.4	28,965	9.8	22,351	13.7	28,848	7.9	72,540	71.6	161,980	30.5
Nation (Prior Year)	1,689,615	63.2	908,604	25.1	1,069,942	16.9	363,039	10.7	466,279	6.1	1,200,700	79.2	4,203,554	9.4

(1) Reporting period is January 1 through December 31.
 (2) TV Viewing and Smoking in Household data are not reported by all contributors.
 (3) Analyses based on one record per child.
 (4) Excludes records with unknown data or errors.
 (5) Rank compares this county's rate to other counties. Rank 1 = best rate.
 (6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 (7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
 (8) Children aged 2 years and older included in the analyses.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Adams	560	9.8 (72)	5.9 (33)	2,344	8.2 (85)	2.5 (58)	10.3 (24)	1,040	16.7 (39)	11.3 (6)	1,662	23.3 (89)
Allen	8,031	9.6 (67)	4.9 (13)	34,121	6.7 (51)	2.4 (54)	11.6 (54)	15,859	16.4 (35)	13.6 (28)	24,597	19.7 (85)
Bartholome	1,540	8.4 (46)	6.0 (38)	5,541	5.9 (29)	1.8 (29)	10.4 (29)	2,340	16.5 (37)	14.9 (47)	3,830	9.6 (22)
Benton	121	10.7 (86)	6.6 (47)	664	4.1 (3)	0.8 (1)	11.6 (54)	352	20.5 (88)	17.6 (79)	496	13.1 (52)
Blackford	280	7.5 (23)	5.4 (26)	1,214	6.7 (51)	3.0 (77)	10.4 (29)	549	12.9 (3)	13.7 (29)	868	10.9 (32)
Boone	562	8.5 (49)	5.9 (33)	2,214	7.1 (67)	4.1 (89)	11.5 (50)	951	15.1 (15)	15.2 (51)	1,503	10.4 (28)
Brown	192	6.8 (12)	10.4 (91)	888	5.2 (14)	1.1 (6)	10.4 (29)	409	14.9 (14)	15.9 (57)	636	2.0 (1)
Carroll	235	9.4 (65)	8.9 (86)	1,107	5.8 (28)	1.4 (13)	11.7 (59)	552	18.7 (71)	16.5 (63)	833	14.9 (66)
Cass	1,043	7.8 (31)	6.6 (47)	4,809	5.1 (10)	2.7 (66)	12.7 (78)	2,379	16.6 (38)	16.8 (67)	3,572	11.6 (39)
Clark	2,082	9.0 (56)	5.6 (28)	8,449	7.3 (73)	1.6 (22)	12.5 (72)	3,568	17.5 (51)	16.7 (66)	5,944	15.6 (68)
Clay	652	5.4 (3)	7.1 (59)	2,977	6.8 (57)	1.2 (9)	13.3 (81)	1,426	18.8 (73)	17.1 (71)	2,097	11.9 (42)

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this county's rate to other counties. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
 - (7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Clinton	838	6.7 (10)	9.3 (88)	3,885	6.7 (51)	1.1 (6)	14.0 (88)	1,865	18.7 (71)	17.6 (79)	2,867	12.8 (47)
Crawford	201	7.0 (14)	5.0 (17)	847	7.9 (83)	1.5 (18)	9.8 (17)	380	18.2 (65)	14.5 (41)	599	11.0 (33)
Daviess	559	10.6 (84)	2.7 (1)	2,405	7.0 (61)	2.7 (66)	14.3 (90)	1,146	17.5 (51)	22.0 (91)	1,682	19.7 (85)
Dearborn	888	7.7 (27)	8.4 (79)	3,538	5.7 (26)	1.4 (13)	10.3 (24)	1,512	17.6 (54)	14.2 (38)	2,510	11.0 (33)
Decatur	552	8.9 (53)	5.4 (26)	2,451	5.5 (21)	1.5 (18)	12.6 (76)	1,158	19.3 (76)	17.6 (79)	1,808	6.7 (6)
DeKalb	887	7.7 (27)	7.8 (73)	3,103	6.5 (42)	2.2 (46)	12.4 (69)	1,321	21.0 (89)	17.3 (74)	2,039	13.6 (55)
Delaware	2,478	10.5 (83)	4.4 (7)	8,820	6.6 (48)	6.1 (92)	10.3 (24)	3,548	15.9 (26)	14.4 (40)	6,015	19.5 (84)
Dubois	736	6.7 (10)	6.7 (49)	3,014	5.6 (23)	2.2 (46)	10.2 (22)	1,310	15.8 (23)	15.1 (49)	2,181	9.1 (20)
Elkhart	5,542	8.1 (38)	7.7 (70)	23,897	5.3 (17)	1.4 (13)	11.6 (54)	11,097	17.6 (54)	16.1 (58)	17,289	6.6 (5)
Fayette	633	7.9 (33)	7.1 (59)	2,438	7.5 (77)	2.0 (41)	10.2 (22)	1,053	15.3 (16)	13.4 (24)	1,643	18.4 (81)
Floyd	1,256	10.1 (77)	5.0 (17)	5,072	8.4 (87)	1.8 (29)	13.9 (87)	2,156	19.6 (80)	18.2 (85)	3,488	12.4 (45)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			(8)	
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Fountain	325	8.3 (44)	7.1 (59)	1,349	5.6 (23)	1.3 (11)	10.0 (20)	604	18.4 (68)	15.6 (54)	949	11.5 (38)
Franklin	294	6.1 (4)	7.1 (59)	1,126	7.0 (61)	1.8 (29)	12.1 (64)	486	20.0 (84)	18.5 (86)	766	21.5 (87)
Fulton	457	8.1 (38)	5.0 (17)	1,971	4.3 (4)	3.2 (83)	12.1 (64)	877	15.7 (21)	17.3 (74)	1,423	15.9 (70)
Gibson	549	7.1 (16)	3.8 (3)	2,166	7.0 (61)	1.8 (29)	11.8 (60)	895	16.2 (33)	17.4 (78)	1,480	13.9 (58)
Grant	1,677	10.2 (80)	5.2 (23)	6,950	7.3 (73)	2.7 (66)	12.4 (69)	3,172	15.8 (23)	15.5 (53)	4,967	22.5 (88)
Greene	498	7.6 (24)	6.2 (42)	2,108	7.2 (72)	1.6 (22)	13.3 (81)	945	19.6 (80)	18.6 (87)	1,511	8.5 (15)
Hamilton	1,902	9.0 (56)	6.8 (51)	8,061	5.9 (29)	2.7 (66)	8.2 (2)	3,471	14.0 (6)	10.1 (2)	5,701	14.0 (59)
Hancock	795	6.5 (8)	7.7 (70)	3,000	8.7 (89)	1.4 (13)	11.8 (60)	1,176	16.4 (35)	13.7 (29)	2,052	12.2 (44)
Harrison	677	9.7 (69)	7.2 (65)	2,747	7.7 (81)	1.9 (35)	11.2 (49)	1,245	17.2 (47)	16.6 (65)	1,941	9.0 (19)
Hendricks	1,321	9.0 (56)	7.0 (57)	4,420	6.5 (42)	3.0 (77)	9.6 (13)	1,726	17.2 (47)	12.3 (15)	2,762	18.4 (81)
Henry	958	9.0 (56)	7.3 (67)	3,999	5.7 (26)	3.3 (85)	9.2 (8)	1,805	17.5 (51)	13.2 (21)	2,925	11.2 (36)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			(8)	
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Howard	2,015	8.4 (46)	5.6 (28)	8,577	7.1 (67)	2.2 (46)	12.1 (64)	3,870	18.1 (62)	15.1 (49)	6,308	24.0 (91)
Huntington	673	7.9 (33)	4.6 (9)	2,797	6.6 (48)	2.6 (62)	9.8 (17)	1,188	14.6 (10)	12.2 (14)	1,998	9.7 (26)
Jackson	988	6.4 (6)	6.7 (49)	4,227	5.6 (23)	1.1 (6)	12.0 (63)	1,891	20.0 (84)	18.0 (84)	3,015	7.0 (7)
Jasper	629	7.9 (33)	7.9 (74)	2,691	5.4 (20)	3.2 (83)	11.6 (54)	1,215	16.9 (43)	14.7 (46)	1,969	17.6 (78)
Jay	423	8.7 (52)	4.0 (4)	1,687	6.5 (42)	1.5 (18)	11.1 (43)	707	17.4 (50)	17.1 (71)	1,179	16.4 (72)
Jefferson	681	11.5 (90)	8.2 (77)	2,687	7.1 (67)	1.6 (22)	13.4 (83)	1,107	21.0 (89)	21.3 (90)	1,863	9.3 (21)
Jennings	789	7.6 (24)	7.7 (70)	3,589	7.6 (80)	3.1 (82)	12.2 (68)	1,748	17.0 (45)	14.9 (47)	2,642	18.8 (83)
Johnson	2,088	7.7 (27)	7.4 (68)	7,759	5.3 (17)	2.6 (62)	7.9 (1)	3,205	15.7 (21)	11.5 (8)	5,457	11.1 (35)
Knox	749	10.3 (81)	4.5 (8)	2,784	6.9 (59)	3.6 (87)	10.3 (24)	1,255	16.0 (27)	14.6 (43)	1,757	5.4 (4)
Kosciusko	1,193	8.5 (49)	6.3 (45)	4,880	6.5 (42)	1.3 (11)	12.8 (79)	2,166	19.3 (76)	17.9 (83)	3,435	8.4 (13)
LaGrange	392	9.7 (69)	5.9 (33)	1,895	6.2 (37)	1.5 (18)	11.1 (43)	923	19.6 (80)	16.5 (63)	1,351	8.8 (16)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Lake	11,935	10.0 (75)	5.6 (28)	48,931	6.3 (39)	2.3 (52)	10.3 (24)	21,700	14.6 (10)	12.1 (12)	34,844	15.7 (69)
LaPorte	2,405	10.3 (81)	5.2 (23)	9,890	6.5 (42)	2.6 (62)	12.1 (64)	4,332	17.7 (57)	13.8 (31)	6,895	12.9 (50)
Lawrence	865	10.1 (77)	5.3 (25)	3,379	7.5 (77)	1.6 (22)	11.0 (40)	1,386	16.9 (43)	13.5 (25)	2,370	8.8 (16)
Madison	2,758	8.4 (46)	5.6 (28)	9,681	9.4 (90)	2.3 (52)	11.5 (50)	3,855	16.3 (34)	14.6 (43)	6,593	23.9 (90)
Marion	27,249	9.7 (69)	6.0 (38)	94,859	6.3 (39)	4.1 (89)	11.1 (43)	39,247	16.1 (29)	14.0 (34)	60,365	17.6 (78)
Marshall	1,004	6.4 (6)	8.0 (75)	4,083	6.9 (59)	1.0 (4)	14.1 (89)	1,804	20.0 (84)	16.9 (68)	2,924	7.1 (8)
Martin	173	9.2 (62)	7.5 (69)	772	5.1 (10)	0.8 (1)	10.8 (37)	333	14.7 (13)	13.5 (25)	547	4.4 (2)
Miami	644	8.5 (49)	8.2 (77)	2,887	6.0 (34)	1.9 (35)	9.4 (10)	1,389	14.4 (8)	11.6 (9)	2,085	11.8 (40)
Monroe	1,814	8.1 (38)	8.7 (83)	7,663	6.7 (51)	1.8 (29)	10.7 (36)	3,392	17.7 (57)	13.9 (32)	5,383	13.3 (53)
Montgomery	812	9.1 (60)	8.0 (75)	3,251	6.4 (41)	2.4 (54)	10.6 (34)	1,433	18.5 (70)	16.1 (58)	2,306	12.8 (47)
Morgan	1,206	8.1 (38)	6.9 (54)	4,462	7.0 (61)	1.7 (27)	11.6 (54)	1,816	19.4 (78)	16.1 (58)	3,103	17.0 (76)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			(8)	
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Newton	177	6.2 (5)	6.8 (51)	759	6.2 (37)	4.5 (91)	13.6 (86)	339	13.6 (5)	12.1 (12)	557	14.2 (62)
Noble	823	7.3 (18)	6.2 (42)	3,285	7.3 (73)	2.7 (66)	12.5 (72)	1,493	16.1 (29)	14.5 (41)	2,198	16.4 (72)
Ohio	20	* *	* *	148	2.0 (1)	2.7 (66)	16.2 (92)	91	* *	* *	115	10.4 (28)
Orange	448	9.6 (67)	4.7 (10)	2,021	7.0 (61)	2.4 (54)	9.6 (13)	988	16.0 (27)	13.3 (23)	1,469	8.2 (12)
Owen	329	7.3 (18)	8.5 (80)	1,480	5.9 (29)	2.5 (58)	12.6 (76)	672	15.8 (23)	17.7 (82)	1,079	9.6 (22)
Parke	235	8.9 (53)	3.4 (2)	955	6.7 (51)	2.0 (41)	10.4 (29)	411	16.1 (29)	12.4 (17)	710	9.6 (22)
Perry	365	11.2 (89)	4.9 (13)	1,308	13.1 (92)	3.7 (88)	15.4 (91)	534	12.7 (2)	9.9 (1)	914	30.7 (92)
Pike	213	11.7 (91)	7.0 (57)	785	11.2 (91)	2.2 (46)	12.5 (72)	293	18.8 (73)	20.8 (89)	519	4.8 (3)
Porter	2,132	6.6 (9)	8.5 (80)	8,399	6.5 (42)	2.2 (46)	10.1 (21)	3,518	15.5 (18)	12.3 (15)	5,899	12.4 (45)
Posey	305	8.9 (53)	4.9 (13)	1,134	5.9 (29)	2.6 (62)	8.2 (2)	429	13.5 (4)	10.5 (4)	763	10.6 (30)
Pulaski	247	6.9 (13)	6.1 (41)	1,090	5.0 (9)	2.2 (46)	11.8 (60)	501	16.8 (41)	12.6 (19)	803	11.2 (36)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Putnam	692	8.1 (38)	8.8 (84)	2,808	5.9 (29)	3.0 (77)	12.4 (69)	1,237	20.2 (87)	16.9 (68)	1,934	8.4 (13)
Randolph	547	9.1 (60)	5.1 (21)	2,115	6.8 (57)	2.1 (43)	9.4 (10)	851	14.5 (9)	12.7 (20)	1,460	14.0 (59)
Ripley	552	9.2 (62)	5.8 (32)	2,416	4.9 (8)	1.9 (35)	10.9 (38)	1,064	16.8 (41)	15.2 (51)	1,746	13.7 (56)
Rush	275	8.0 (36)	6.9 (54)	1,086	5.5 (21)	3.5 (86)	9.1 (6)	472	16.1 (29)	14.0 (34)	744	12.8 (47)
Scott	656	9.9 (73)	4.1 (5)	2,906	6.0 (34)	3.0 (77)	11.5 (50)	1,364	15.6 (19)	16.3 (61)	2,059	13.0 (51)
Shelby	897	7.0 (14)	6.2 (42)	3,394	7.0 (61)	1.0 (4)	13.4 (83)	1,399	16.7 (39)	17.3 (74)	2,334	14.1 (61)
Spencer	270	7.8 (31)	4.1 (5)	1,104	7.5 (77)	0.9 (3)	13.4 (83)	493	19.7 (83)	18.9 (88)	812	8.1 (11)
St Joseph	5,595	9.5 (66)	7.1 (59)	21,762	5.1 (10)	2.9 (74)	9.7 (16)	9,711	14.6 (10)	11.4 (7)	14,413	14.8 (65)
Starke	534	7.3 (18)	9.4 (89)	2,427	5.1 (10)	1.2 (9)	13.0 (80)	1,161	18.3 (66)	16.4 (62)	1,895	15.0 (67)
Steuben	751	8.1 (38)	8.8 (84)	3,847	4.4 (5)	2.9 (74)	10.6 (34)	2,007	17.3 (49)	14.6 (43)	2,859	10.2 (27)
Sullivan	435	7.1 (16)	6.0 (38)	1,571	8.3 (86)	1.7 (27)	12.5 (72)	640	19.5 (79)	16.9 (68)	1,084	12.0 (43)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			(8)	
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Switzerland	121	2.5 (1)	9.1 (87)	561	4.8 (6)	2.9 (74)	10.5 (33)	257	18.3 (66)	14.0 (34)	414	11.8 (40)
Tippecanoe	3,584	7.4 (22)	8.5 (80)	14,864	4.8 (6)	2.5 (58)	11.0 (40)	6,652	17.1 (46)	13.9 (32)	10,503	13.7 (56)
Tipton	172	9.3 (64)	4.7 (10)	797	5.3 (17)	3.0 (77)	9.2 (8)	395	12.4 (1)	13.2 (21)	598	7.5 (10)
Union	132	8.3 (44)	9.8 (90)	586	3.9 (2)	1.4 (13)	9.4 (10)	279	21.9 (91)	11.8 (10)	415	17.6 (78)
Vanderburg	3,381	10.6 (84)	4.7 (10)	12,059	7.1 (67)	1.9 (35)	9.9 (19)	4,642	17.8 (61)	14.2 (38)	8,180	8.8 (16)
Vermillion	286	10.1 (77)	5.9 (33)	1,191	7.7 (81)	1.9 (35)	11.1 (43)	517	17.6 (54)	17.2 (73)	867	16.4 (72)
Vigo	2,247	7.7 (27)	7.1 (59)	7,937	6.7 (51)	1.9 (35)	10.9 (38)	3,110	17.7 (57)	15.8 (56)	5,427	10.7 (31)
Wabash	607	7.6 (24)	5.9 (33)	2,842	6.0 (34)	1.8 (29)	9.1 (6)	1,375	15.3 (16)	11.0 (5)	2,047	7.3 (9)
Warren	175	5.1 (2)	6.9 (54)	794	5.2 (14)	2.5 (58)	9.6 (13)	386	18.1 (62)	13.5 (25)	557	13.5 (54)
Warrick	740	9.9 (73)	7.2 (65)	2,862	8.6 (88)	2.8 (73)	11.1 (43)	1,174	14.3 (7)	12.4 (17)	1,973	14.7 (63)
Washington	470	10.0 (75)	4.9 (13)	2,202	7.3 (73)	2.1 (43)	11.0 (40)	1,022	19.2 (75)	15.6 (54)	1,604	9.6 (22)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			(8)	
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Wayne	1,687	8.0 (36)	6.4 (46)	6,729	6.6 (48)	2.1 (43)	11.1 (43)	2,893	17.7 (57)	14.1 (37)	4,612	16.5 (75)
Wells	422	10.9 (87)	5.0 (17)	1,733	7.1 (67)	1.6 (22)	8.9 (4)	764	18.1 (62)	11.9 (11)	1,248	14.7 (63)
White	603	7.3 (18)	6.8 (51)	2,529	5.2 (14)	2.4 (54)	8.9 (4)	1,143	15.6 (19)	10.1 (2)	1,832	17.5 (77)
Whitley	565	11.0 (88)	5.1 (21)	2,439	7.9 (83)	2.7 (66)	11.5 (50)	1,115	18.4 (68)	17.3 (74)	1,755	16.3 (71)
Unknown Counties	2	* *	* *	5	* *	* *	* *	2	* *	* *	3	* *
Indiana	132,498	9.0	6.3	522,056	6.4	2.6	11.1	227,729	16.6	14.3	361,426	14.6

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
County	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
Adams	624	78.2 (4)	468	22.2 (12)	453	9.1 (31)	457	17.5 (12)	463	10.8 (12)	1,117	67.2 (78)	1,308	37.7 (33)
Allen	8,757	60.8 (54)	6,627	17.7 (30)	6,837	8.6 (36)	6,239	11.3 (59)	6,577	7.7 (45)	16,131	73.1 (59)	18,686	32.3 (25)
Bartholome	1,717	67.0 (32)	1,228	16.7 (38)	1,130	7.6 (52)	1,184	12.8 (46)	1,223	8.7 (29)	2,487	85.0 (9)	3,387	47.3 (56)
Benton	135	59.3 (66)	109	12.8 (68)	114	9.6 (25)	102	19.6 (8)	108	5.6 (68)	376	68.1 (77)	347	44.7 (45)
Blackford	300	70.0 (24)	236	15.7 (46)	234	6.4 (63)	239	16.3 (16)	236	11.0 (11)	585	74.7 (45)	639	50.7 (64)
Boone	645	57.4 (71)	453	19.9 (19)	462	9.3 (28)	439	13.4 (42)	449	10.0 (15)	1,009	73.5 (55)	1,316	46.8 (53)
Brown	221	78.7 (2)	172	32.6 (1)	175	18.9 (1)	163	27.6 (2)	172	8.7 (29)	439	82.9 (13)	484	47.7 (57)
Carroll	256	70.3 (19)	214	20.1 (17)	209	9.6 (25)	202	15.8 (18)	213	8.5 (32)	580	87.2 (5)	586	19.1 (9)
Cass	1,132	59.5 (63)	942	16.1 (44)	879	7.8 (50)	901	7.7 (85)	940	4.1 (87)	2,562	63.7 (84)	2,578	19.5 (10)
Clark	2,379	66.1 (37)	1,795	21.1 (16)	1,746	11.4 (13)	1,679	15.6 (19)	1,779	10.3 (14)	3,840	73.4 (57)	4,820	39.4 (36)
Clay	712	61.7 (52)	571	15.1 (49)	566	6.2 (66)	555	10.3 (68)	569	7.4 (48)	1,528	75.4 (42)	1,643	52.1 (73)
Clinton	916	71.4 (14)	796	16.8 (36)	731	9.2 (29)	749	8.7 (83)	792	5.2 (74)	2,013	66.1 (81)	2,130	14.0 (4)
Crawford	221	64.7 (43)	191	17.8 (29)	180	11.1 (14)	180	17.8 (11)	190	10.5 (13)	409	80.2 (22)	481	54.1 (80)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
County	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
Daviess	628	56.7 (74)	438	12.1 (72)	466	5.8 (75)	400	5.0 (90)	436	4.6 (84)	1,239	69.8 (70)	1,368	36.3 (31)
Dearborn	995	59.9 (60)	754	12.5 (70)	715	5.5 (77)	717	12.6 (49)	751	7.6 (46)	1,600	81.6 (19)	2,055	57.2 (88)
Decatur	597	64.3 (45)	502	13.3 (62)	457	5.5 (77)	485	11.8 (55)	497	7.8 (41)	1,238	78.8 (26)	1,347	56.9 (87)
DeKalb	985	78.6 (3)	669	17.6 (32)	657	8.2 (41)	659	16.4 (15)	663	9.7 (20)	1,407	76.1 (37)	1,796	52.2 (75)
Delaware	2,601	60.0 (58)	1,739	16.2 (43)	1,661	8.7 (34)	1,719	11.5 (58)	1,729	5.6 (68)	3,813	72.8 (60)	5,001	23.7 (13)
Dubois	817	74.3 (9)	702	23.5 (9)	620	11.8 (9)	674	14.4 (30)	700	8.3 (36)	1,405	76.9 (33)	1,657	37.6 (32)
Elkhart	6,012	73.6 (10)	4,587	26.3 (6)	4,597	12.4 (7)	4,289	13.3 (43)	4,545	8.5 (32)	11,918	74.6 (46)	13,142	8.7 (2)
Fayette	685	56.9 (72)	475	13.5 (60)	499	6.6 (62)	455	9.0 (80)	475	4.0 (88)	1,120	77.1 (31)	1,417	52.9 (79)
Floyd	1,449	64.9 (42)	1,035	19.1 (24)	1,018	8.7 (34)	1,001	12.9 (45)	1,024	7.8 (41)	2,288	73.5 (55)	2,959	46.3 (52)
Fountain	363	62.8 (48)	273	15.0 (50)	267	8.6 (36)	272	7.7 (85)	272	5.1 (75)	656	72.3 (64)	782	50.8 (66)
Franklin	323	60.1 (57)	224	11.6 (75)	220	8.2 (41)	205	11.2 (61)	223	4.9 (78)	525	86.1 (7)	672	55.5 (83)
Fulton	503	66.2 (36)	424	14.6 (52)	409	6.4 (63)	392	13.5 (40)	421	8.1 (37)	922	76.6 (34)	1,118	27.6 (18)
Gibson	632	51.1 (88)	501	11.2 (80)	410	5.6 (76)	475	10.7 (65)	501	6.8 (56)	937	73.6 (54)	1,299	48.5 (58)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
County	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
Grant	1,816	52.3 (85)	1,469	10.6 (83)	1,391	5.2 (81)	1,384	8.5 (84)	1,462	5.5 (71)	3,417	65.7 (83)	3,804	26.6 (17)
Greene	529	58.2 (70)	440	7.3 (90)	420	4.5 (89)	419	9.1 (79)	440	4.8 (80)	997	94.2 (1)	1,176	39.0 (35)
Hamilton	2,255	74.8 (7)	1,821	27.4 (3)	1,658	14.2 (5)	1,744	20.7 (6)	1,807	13.6 (5)	3,670	84.7 (12)	4,677	21.4 (11)
Hancock	896	70.6 (17)	616	21.4 (14)	622	10.1 (22)	607	20.8 (5)	612	14.2 (4)	1,269	73.8 (53)	1,799	16.5 (5)
Harrison	746	70.2 (21)	598	13.7 (59)	552	6.7 (60)	560	16.3 (16)	598	8.9 (28)	1,326	82.3 (15)	1,578	44.0 (43)
Hendricks	1,493	70.1 (22)	864	22.2 (12)	876	10.3 (21)	854	19.0 (9)	856	12.9 (6)	1,835	80.5 (20)	2,798	38.8 (34)
Henry	1,031	66.9 (33)	823	16.3 (42)	817	8.2 (41)	771	14.5 (28)	819	9.9 (16)	1,923	66.6 (80)	2,174	51.5 (70)
Howard	2,231	53.6 (82)	1,932	11.9 (73)	1,792	6.2 (66)	1,771	11.3 (59)	1,928	7.3 (50)	4,152	74.3 (51)	4,704	35.9 (30)
Huntington	747	64.7 (43)	572	14.0 (58)	559	8.1 (45)	543	14.0 (33)	571	9.5 (21)	1,273	75.0 (43)	1,569	49.8 (62)
Jackson	1,093	67.6 (28)	867	14.6 (52)	840	7.0 (56)	823	11.7 (56)	865	7.3 (50)	2,034	78.2 (29)	2,291	42.4 (41)
Jasper	700	60.4 (56)	582	17.9 (27)	540	9.6 (25)	549	18.4 (10)	579	11.9 (8)	1,305	53.4 (89)	1,544	46.8 (53)
Jay	476	65.5 (39)	356	17.7 (30)	329	8.2 (41)	338	13.9 (34)	352	9.1 (26)	756	59.0 (88)	968	44.8 (46)
Jefferson	760	66.3 (35)	540	16.7 (38)	505	10.9 (15)	536	14.9 (23)	535	9.5 (21)	1,202	76.5 (35)	1,543	28.6 (21)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
County	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
Jennings	848	61.0 (53)	666	13.2 (64)	655	6.0 (71)	602	11.6 (57)	665	8.0 (39)	1,877	78.3 (28)	1,923	50.0 (63)
Johnson	2,276	69.7 (25)	1,826	19.2 (23)	1,539	10.0 (24)	1,723	12.4 (50)	1,816	6.6 (58)	3,362	76.1 (37)	4,581	31.3 (24)
Knox	816	50.7 (89)	493	12.6 (69)	561	7.0 (56)	479	9.6 (73)	493	5.9 (66)	1,329	92.6 (2)	1,694	34.5 (28)
Kosciusko	1,340	70.3 (19)	1,017	23.5 (9)	1,017	10.6 (17)	963	15.5 (21)	1,010	8.1 (37)	2,316	72.5 (61)	2,753	33.3 (26)
LaGrange	449	72.4 (12)	378	21.4 (14)	331	14.5 (4)	356	19.7 (7)	376	11.7 (9)	973	84.8 (11)	1,077	42.5 (42)
Lake	13,170	66.7 (34)	10,570	15.3 (48)	9,548	8.1 (45)	9,953	7.1 (88)	10,523	4.4 (86)	23,150	63.0 (86)	27,777	23.5 (12)
LaPorte	2,705	60.0 (58)	2,170	19.5 (21)	1,854	12.2 (8)	2,012	8.9 (82)	2,150	5.5 (71)	4,308	51.6 (90)	5,609	34.6 (29)
Lawrence	930	65.2 (41)	733	11.3 (77)	677	5.9 (73)	692	10.8 (63)	732	6.1 (63)	1,475	79.3 (24)	1,865	46.0 (50)
Madison	2,982	56.4 (75)	2,066	17.6 (32)	1,877	10.1 (22)	1,945	13.6 (36)	2,054	9.4 (24)	4,091	68.8 (73)	5,802	45.4 (48)
Marion	29,389	62.0 (51)	18,365	23.6 (8)	19,894	11.7 (10)	17,653	9.6 (73)	18,154	6.0 (64)	32,140	66.7 (79)	53,829	24.1 (15)
Marshall	1,108	71.6 (13)	884	18.3 (26)	804	9.2 (29)	850	14.4 (30)	880	9.8 (18)	1,930	75.0 (43)	2,356	8.3 (1)
Martin	190	53.7 (81)	163	11.7 (74)	149	6.7 (60)	160	10.6 (67)	163	6.7 (57)	358	90.5 (3)	417	33.6 (27)
Miami	684	59.6 (62)	590	12.9 (66)	571	4.7 (86)	545	13.6 (36)	589	7.6 (46)	1,463	72.5 (61)	1,566	46.9 (55)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
County	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
Monroe	2,060	77.8 (5)	1,660	29.5 (2)	1,517	18.5 (2)	1,574	24.7 (3)	1,637	17.5 (1)	3,580	86.6 (6)	4,327	17.0 (8)
Montgomery	909	66.0 (38)	685	14.2 (56)	669	4.9 (83)	665	9.9 (71)	681	5.6 (68)	1,531	63.3 (85)	1,870	46.0 (50)
Morgan	1,332	67.6 (28)	997	15.0 (50)	835	6.3 (65)	946	14.2 (32)	994	8.4 (35)	1,935	66.0 (82)	2,652	56.0 (85)
Newton	201	58.7 (67)	169	17.2 (34)	163	8.0 (49)	147	13.6 (36)	169	11.2 (10)	372	29.0 (91)	447	55.7 (84)
Noble	901	70.6 (17)	601	22.3 (11)	686	10.5 (19)	576	17.5 (12)	597	12.4 (7)	1,585	68.7 (75)	1,938	29.2 (23)
Ohio	23	* *	22	* *	22	* *	19	* *	22	* *	91	* *	71	* *
Orange	481	70.7 (16)	415	16.9 (35)	360	7.2 (54)	391	15.1 (22)	413	5.1 (75)	1,053	71.3 (67)	1,097	26.1 (16)
Owen	374	67.6 (28)	299	16.4 (41)	292	10.6 (17)	277	13.7 (35)	297	4.7 (82)	714	78.4 (27)	823	52.5 (77)
Parke	258	58.5 (68)	199	7.5 (89)	190	5.3 (80)	188	9.0 (80)	198	4.0 (88)	437	74.4 (48)	549	54.1 (80)
Perry	397	49.6 (91)	307	7.2 (91)	235	3.0 (91)	324	6.5 (89)	306	3.3 (90)	576	73.4 (57)	800	60.5 (90)
Pike	233	54.5 (79)	174	11.5 (76)	148	6.1 (69)	180	9.4 (78)	171	2.9 (91)	310	71.3 (67)	446	60.5 (90)
Porter	2,393	70.1 (22)	1,867	18.6 (25)	1,639	10.7 (16)	1,772	14.5 (28)	1,860	9.1 (26)	3,755	75.6 (41)	4,913	28.5 (19)
Posey	355	52.7 (83)	285	9.1 (88)	220	5.9 (73)	270	9.6 (73)	282	6.4 (59)	460	74.1 (52)	669	49.0 (59)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
County	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
Pulaski	266	59.4 (64)	235	10.2 (85)	210	6.2 (66)	211	14.7 (25)	235	7.2 (52)	531	74.4 (48)	609	40.7 (39)
Putnam	749	62.6 (49)	550	12.4 (71)	574	6.8 (58)	522	12.3 (51)	547	4.9 (78)	1,299	74.4 (48)	1,604	54.6 (82)
Randolph	616	59.9 (60)	450	19.3 (22)	435	8.5 (38)	424	15.6 (19)	448	9.8 (18)	895	59.6 (87)	1,199	40.6 (38)
Ripley	605	56.0 (77)	480	9.8 (87)	481	6.0 (71)	458	7.6 (87)	479	4.8 (80)	1,124	79.0 (25)	1,316	56.8 (86)
Rush	298	54.0 (80)	218	13.3 (62)	225	4.0 (90)	199	12.1 (53)	216	4.6 (84)	505	80.4 (21)	622	49.7 (61)
Scott	740	51.9 (86)	547	11.3 (77)	538	4.8 (85)	528	9.5 (76)	544	7.2 (52)	1,444	77.1 (31)	1,618	52.8 (78)
Shelby	1,009	63.4 (47)	765	16.5 (40)	658	8.1 (45)	735	13.5 (40)	761	8.5 (32)	1,489	68.8 (73)	1,970	49.5 (60)
Spencer	315	51.4 (87)	257	14.4 (54)	237	6.8 (58)	253	9.5 (76)	255	5.5 (71)	539	85.0 (9)	672	52.2 (75)
St Joseph	6,136	73.1 (11)	4,081	26.5 (5)	4,313	13.7 (6)	3,924	14.8 (24)	4,043	9.2 (25)	10,341	76.1 (37)	12,650	16.7 (6)
Starke	577	65.5 (39)	488	16.8 (36)	481	8.3 (39)	451	14.6 (27)	483	9.9 (16)	1,258	77.8 (30)	1,302	51.2 (68)
Steuben	835	78.8 (1)	688	20.1 (17)	701	8.3 (39)	635	16.7 (14)	684	9.5 (21)	2,131	79.5 (23)	2,085	51.0 (67)
Sullivan	485	68.7 (27)	321	10.0 (86)	274	4.7 (86)	307	13.0 (44)	321	6.2 (62)	676	68.5 (76)	940	51.7 (71)
Switzerlan	140	50.7 (89)	114	10.5 (84)	106	10.4 (20)	109	12.8 (46)	112	7.1 (54)	267	85.8 (8)	310	60.3 (89)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
County	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months	Number	% Breastfed At Least 3 Months	Number	% Breastfed At Least 6 Months	Number	%		
	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)		
Tippecanoe	3,951	74.5 (8)	2,970	27.0 (4)	2,986	16.3 (3)	2,764	21.2 (4)	2,942	16.3 (3)	7,014	74.5 (47)	8,433	28.5 (19)
Tipton	191	62.3 (50)	149	13.4 (61)	151	7.3 (53)	143	14.7 (25)	148	7.4 (48)	415	81.7 (16)	449	28.7 (22)
Union	156	56.4 (75)	112	17.9 (27)	114	8.8 (33)	99	* *	112	8.0 (39)	294	87.4 (4)	335	51.3 (69)
Vanderburg	3,777	55.1 (78)	2,907	10.9 (82)	2,474	6.1 (69)	2,730	10.3 (68)	2,899	6.4 (59)	4,971	72.4 (63)	7,245	11.1 (3)
Vermillion	301	52.5 (84)	232	12.9 (66)	240	5.0 (82)	219	11.0 (62)	232	7.8 (41)	551	71.3 (67)	653	52.1 (73)
Vigo	2,441	58.3 (69)	1,693	11.1 (81)	1,609	5.5 (77)	1,623	9.9 (71)	1,690	5.9 (66)	3,305	69.5 (71)	4,689	50.7 (64)
Wabash	647	64.0 (46)	562	14.2 (56)	533	4.9 (83)	509	10.8 (63)	561	7.8 (41)	1,467	71.4 (66)	1,497	24.0 (14)
Warren	204	60.8 (54)	150	11.3 (77)	153	4.6 (88)	150	10.7 (65)	149	4.7 (82)	405	71.6 (65)	447	51.7 (71)
Warrick	826	56.8 (73)	668	13.2 (64)	555	7.2 (54)	665	12.8 (46)	667	6.3 (61)	1,257	76.5 (35)	1,649	45.7 (49)
Washington	540	69.6 (26)	463	14.3 (55)	433	8.1 (45)	450	12.2 (52)	460	5.0 (77)	1,108	75.7 (40)	1,194	39.7 (37)
Wayne	1,867	59.4 (64)	1,396	16.0 (45)	1,411	9.0 (32)	1,342	10.1 (70)	1,384	6.0 (64)	3,084	81.7 (16)	3,765	41.3 (40)
Wells	465	71.2 (15)	368	15.5 (47)	357	7.8 (50)	354	13.6 (36)	368	8.7 (29)	812	69.1 (72)	961	45.3 (47)
White	659	67.2 (31)	557	19.9 (19)	488	11.7 (10)	528	12.1 (53)	554	6.9 (55)	1,232	82.8 (14)	1,346	16.9 (7)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this county's rate to other counties. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7B
2009 - 2011 Pediatric Nutrition Surveillance (1)
Indiana

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by County (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
County	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
Whitley	621	75.4 (6)	468	25.9 (7)	473	11.6 (12)	456	28.3 (1)	466	17.2 (2)	1,198	81.7 (16)	1,401	44.5 (44)
Unknown Counties	5	* *	3	* *	2	* *	4	* *	3	* *	2	* *	7	* *
Indiana	145,604	64.6	107,098	18.8	104,278	9.7	101,785	12.2	106,367	7.6	232,088	72.0	296,462	31.1

(1) Reporting period is January 1 through December 31.
 (2) TV Viewing and Smoking in Household data are not reported by all contributors.
 (3) Analyses based on one record per child.
 (4) Excludes records with unknown data or errors.
 (5) Rank compares this county's rate to other counties. Rank 1 = best rate.
 (6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
 (7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
 (8) Children aged 2 years and older included in the analyses.
 * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

